

UZEDY[®]
(risperidone)
extended-release
injectable suspension

50 mg 75 mg 100 mg 125 mg
150 mg 200 mg 250 mg

SEE WHAT FEWER BIPOLAR SYMPTOMS* MIGHT DO FOR

With bipolar I disorder,
you may have highs and lows

A DIFFERENT TREATMENT OPTION MAY BE A FIT FOR YOU

People featured are not actual
patients, caregivers, or doctors.

*In the clinical trial, all participants first received another long-acting risperidone injection. About 60% of them (303 out of 501 people) were judged to be stable and then randomized to either the same dose of risperidone long-acting injection or placebo and monitored for relapse. Time to relapse was delayed in patients receiving risperidone long-acting injection, compared with patients receiving placebo. The majority of relapses were due to manic rather than depressive symptoms.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY[®]?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at [UZEDY.com](#).

teva



MISSING DAILY PILLS CAN SET TREATMENT BACK AND INCREASE RELAPSE RISK

A LONGER-LASTING OPTION MAY HELP KEEP YOU STEADY



A relapse is a sudden return to a severe mood episode. This can be an episode of either mania or depression.



It can be common for people with bipolar I disorder to not take their daily pills as prescribed. Missing doses can lead to mood episodes, relapses, and even hospital visits.



Long-acting injectable (LAI) treatments don't have to be taken every day and provide long-lasting symptom control.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Do not receive UZEDY® if you are allergic to risperidone, paliperidone, or any of its components.

UZEDY may cause serious side effects, including:

- **Stroke in elderly people (cerebrovascular problems) that can lead to death.**
- **Neuroleptic Malignant Syndrome (NMS). NMS is a rare but very serious problem that can lead to death. Seek medical attention right away** if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in your blood pressure.
- **Uncontrolled facial or body movements (tardive dyskinesia)** may not go away, even if you stop receiving UZEDY. Tardive dyskinesia may also start after you stop receiving UZEDY.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including **Boxed WARNING, at [UZEDY.com](#).**

WITH UZEDY, YOU HAVE OPTIONS

UZEDY® is an LAI treatment for bipolar I disorder that can help prevent relapses.* UZEDY is given as an injection by a healthcare provider, such as a doctor or nurse. UZEDY is injected just below the skin. You and your doctor get to decide if you'd rather have UZEDY injected in your arm or abdomen.



YOU DON'T HAVE TO TAKE A PILL EVERY DAY

With UZEDY, you get an injection once a month.

A STREAMLINED START

No extra risperidone pills or additional starter doses are needed once you begin UZEDY. It's absorbed rapidly after it's injected and then slowly releases medicine into your body over time.

UZEDY MAY PROVIDE DELAYED TIME TO RELAPSE



In the clinical trial, all participants first received another long-acting risperidone injection. About 60% of them (303 out of 501 people) were judged to be stable and then randomized to either the same dose of risperidone long-acting injection or placebo and monitored for relapse. Time to relapse was delayed in patients receiving risperidone long-acting injection, compared with patients receiving placebo.[†]

*The efficacy of UZEDY is based on another risperidone long-acting injection that was administered into the muscle. UZEDY works similarly but is injected just below the skin.

[†]The majority of relapses were due to manic rather than depressive symptoms.

IMPORTANT SAFETY INFORMATION (CONTINUED)


- **Problems with your metabolism** that may include high blood sugar (hyperglycemia), diabetes mellitus, changes in the fat levels in your blood (dyslipidemia), and weight gain. Extremely high blood sugar can lead to coma or death. If you have diabetes or are at risk for diabetes (e.g., obesity, family history of diabetes), your healthcare provider should check your blood sugar before you start and during treatment with UZEDY. Call your healthcare provider if you have symptoms of high blood sugar including: feeling very thirsty, hungry, sick to your stomach, weak or tired, or confused; needing to urinate more than usual; or your breath smells fruity.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including **Boxed WARNING**, at [UZEDY.com](#).



HOW TREATMENT WITH UZEDY® IS DIFFERENT

UZEDY HAS A SHORTER NEEDLE THAN OTHER LAIs FOR BIPOLAR I DISORDER*



IN A SURVEY, 7 OUT OF 10 PEOPLE FELT UZEDY WAS A BETTER INJECTION EXPERIENCE THAN OTHER LAIs THEY'D TAKEN IN THE PAST[‡]

89% OF PEOPLE (56 OUT OF 63) IN A SURVEY SAID RECEIVING UZEDY WAS EASY[‡]

*UZEDY is supplied as a prefilled syringe designed to provide 0.14 mL to 0.28 mL of medication, depending on the dose, with a 5/8-inch needle. No conclusions can be made between needle length and safety and effectiveness as this has not been studied.
[†]Shortest needle length; additional needle lengths for gluteal injection and/or patients with body weight over certain thresholds.
[‡]Data were collected from 63 patients, 24 physicians, and 25 nurses in a prospective, cross-sectional companion survey assessing the perceptions regarding ease of use and satisfaction with UZEDY. The survey was administered after a minimum of 2 experiences prescribing, administering, or receiving UZEDY.

IMPORTANT SAFETY INFORMATION (CONTINUED)

- **High levels of prolactin in your blood.** UZEDY may cause a rise in the blood levels of a hormone called prolactin that may cause side effects including missed menstrual periods, decreased fertility in women, leakage of milk from the breasts, development of breasts in men, or problems with erection.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including **Boxed WARNING**, at [UZEDY.com](#).

All trademarks are the property of their respective owners.

START THE CONVERSATION WITH YOUR DOCTOR

Because bipolar I disorder affects everyone differently, talking openly with your doctor can help them better understand what you're experiencing. Sharing your thoughts and questions can make it easier to find a treatment plan that works for you.



TEVA OFFERS SUPPORT TO HELP YOU PLAN FOR STARTING AND STAYING ON UZEDY®

We help streamline your start to treatment by helping you, or a caregiver, figure out some of the details.

SUPPORT FROM TEVA CAN:

- ✓ Confirm prescription coverage and pharmacy options based on eligibility and coverage
- ✓ Explain Medicare and Medicaid coverage options
- ✓ See what financial assistance options may be available to you
- ✓ Help you understand prior authorization requirements
- ✓ Work with you to find an injection site near your home or work
- ✓ Provide over-the-phone nurse support to answer questions and help you stay on UZEDY

TALK TO YOUR DOCTOR OR VISIT [UZEDY.COM](https://uzedy.com) TO LEARN MORE

IMPORTANT SAFETY INFORMATION (CONTINUED)

- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- **Falls.** Antipsychotic medicines like UZEDY may cause drowsiness or dizziness when you are standing, which could increase your risk for falls and related injuries.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at [UZEDY.com](https://uzedy.com).



APPROVED USE

UZEDY® (risperidone) extended-release injectable suspension is a prescription medicine used in adults for the maintenance treatment of bipolar I disorder as monotherapy or as adjunctive therapy to lithium or valproate.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Do not receive UZEDY if you are allergic to risperidone, paliperidone, or any of its components.

UZEDY may cause serious side effects, including:

- **Stroke in elderly people (cerebrovascular problems) that can lead to death.**
- **Neuroleptic Malignant Syndrome (NMS). NMS is a rare but very serious problem that can lead to death. Seek medical attention right away** if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in your blood pressure.
- **Uncontrolled facial or body movements (tardive dyskinesia)** may not go away, even if you stop receiving UZEDY. Tardive dyskinesia may also start after you stop receiving UZEDY.
- **Problems with your metabolism** that may include high blood sugar (hyperglycemia),

diabetes mellitus, changes in the fat levels in your blood (dyslipidemia), and weight gain. Extremely high blood sugar can lead to coma or death. If you have diabetes or are at risk for diabetes (e.g., obesity, family history of diabetes), your healthcare provider should check your blood sugar before you start and during treatment with UZEDY. Call your healthcare provider if you have symptoms of high blood sugar including: feeling very thirsty, hungry, sick to your stomach, weak or tired, or confused; needing to urinate more than usual; or your breath smells fruity.

- **High levels of prolactin in your blood.** UZEDY may cause a rise in the blood levels of a hormone called prolactin that may cause side effects including missed menstrual periods, decreased fertility in women, leakage of milk from the breasts, development of breasts in men, or problems with erection.
- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- **Falls.** Antipsychotic medicines like UZEDY may cause drowsiness or dizziness when you are standing, which could increase your risk for falls and related injuries.
- **Low white blood cell count.**
- **Problems thinking clearly and moving your body.** Do not drive, operate machinery, or do other dangerous activities until you know how UZEDY affects you.

Please see continued Important Safety Information on page 7 and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at UZEDY.com.

- **Seizures (convulsions).**
- **Difficulty swallowing that can cause food or liquid to get into your lungs.**
- **Prolonged or painful erection lasting more than 4 hours.** Call your healthcare provider or go to your nearest emergency room right away if you have an erection that lasts more than 4 hours.
- **Problems with control of your body temperature (too high or too low). Avoid getting overheated or dehydrated.**

The most common side effects of risperidone in patients with

- **Bipolar disorder** were weight increased (5% in monotherapy trial) and slow movements, stiffness, shaking ($\geq 10\%$ in adjunctive therapy trial).
- **Injection site reactions** including a lump or itching were reported with UZEDY®.

These are not all the possible side effects of UZEDY. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. For more information, ask your healthcare provider or pharmacist.

Do not drink alcohol during treatment with UZEDY.

Before receiving UZEDY, tell your healthcare provider about all your medical conditions, including if you:

- have had Neuroleptic Malignant Syndrome.
- have or have had uncontrolled movements of your tongue, face, mouth, or jaw (tardive dyskinesia).
- have diabetes or have a family history of diabetes.
- have had dizziness or fainting or are being

treated for high blood pressure.

- have had a low white blood cell count.
- have or have had seizures or epilepsy.
- are pregnant or plan to become pregnant during treatment with UZEDY. It is not known if UZEDY will harm your unborn baby. Use of UZEDY during the third trimester of pregnancy may cause side effects in the newborn infant, including agitation, abnormal muscle tone, tremor, drowsiness, difficulty feeding, and difficulty breathing. Seek medical attention if you notice these signs. If you become pregnant during treatment with UZEDY, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics, or call [1-866-961-2388](tel:1-866-961-2388) or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.
- are breastfeeding or plan to breastfeed. If you are receiving UZEDY and are breastfeeding, monitor your infant for sleepiness, inadequate weight gain, jitteriness, tremors, and abnormal muscle movements. Seek medical care if you notice these signs.
- have or have had kidney or liver problems.

Tell your healthcare provider about all the medicines you take or plan to take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. UZEDY and other medicines may affect each other.

You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

For more information about UZEDY, see the full [Prescribing Information](#), including **Boxed WARNING, or talk to your healthcare provider.**

UZEDY
(risperidone)
extended-release
injectable suspension

50 mg, 75 mg, 100 mg, 125 mg

150 mg, 200 mg, 250 mg



GOT THIS

Learn more about managing your bipolar I disorder with UZEDY® at [UZEDY.com](https://www.UZEDY.com)

People featured are not actual patients, caregivers, or doctors.

APPROVED USE

UZEDY (risperidone) extended-release injectable suspension is a prescription medicine used in adults for the maintenance treatment of bipolar I disorder as monotherapy or as adjunctive therapy to lithium or valproate.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Please see additional Important Safety Information on previous pages and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at [UZEDY.com](https://www.UZEDY.com).

teva

© 2026 Teva Neuroscience, Inc.
RIS-41782 January 2026